arXiv:math/9908140v1 [math.CA] 27 Aug 1999

A note on the q-derivative operator

J. Koekoek R. Koekoek

Abstract

The following statement is proved. If the q-derivative operator $\mathcal{D}_q$ is defined by

$$\mathcal{D}_q f(x) := \begin{cases} \dfrac{f(x)-f(qx)}{(1-q)x}, & x \neq 0 \\ \\ f'(0), & x = 0 \end{cases}$$

for functions f which are differentiable at $x = 0$, then we have for every positive integer n

$$\left(\mathcal{D}_q^n f\right)(0) = \lim_{x\to 0} \mathcal{D}_q^n f(x) = \frac{f^{(n)}(0)}{n!}\frac{(q;q)_n}{(1-q)^n}$$

for every function f whose nth derivative at $x = 0$ exists.

We give a proof in both the real variable and the complex variable case.

Usually the q-derivative operator $\mathcal{D}_q$ is defined by

$$\mathcal{D}_q f(x) = \frac{f(x)-f(qx)}{(1-q)x}$$

where q is fixed and $0 < q < 1$. However, this definition is not valid for $x = 0$. On the other hand, the hypothesis $0 < q < 1$ can be weakened. So we define

$$\mathcal{D}_q f(x) := \begin{cases} \dfrac{f(x)-f(qx)}{(1-q)x}, & x \neq 0 \\ \\ f'(0), & x = 0 \end{cases} \tag{1}$$

for functions f which are differentiable at $x = 0$, where q is a fixed real number not equal to 1.

Note that the existence of $f'(0)$ implies that the domain of f contains the point zero in its interior.

Further we define $\mathcal{D}_q^n f := \mathcal{D}_q\left(\mathcal{D}_q^{n-1} f\right)$ for $n = 1, 2, 3, \ldots$, where $\mathcal{D}_q^0$ denotes the identity operator.

In this paper we prove the following theorem.

Theorem. *Let n be a positive integer and let f be a function for which $f^{(n)}(0)$ exists. Then we have*

$$\left(\mathcal{D}_q^n f\right)(0) = \lim_{x\to 0} \mathcal{D}_q^n f(x) = \frac{f^{(n)}(0)}{n!}\frac{(q;q)_n}{(1-q)^n}.$$

The real variable case.

Let f be a function of a real variable x and let the domain of f contain the interval $(-\rho,\rho)$ for some $\rho > 0$.

We use the following version of l'Hospital's rule (see for instance section 84 in [1]) :

Let F and G be functions whose nth ($n \geq 1$) derivatives at $x = 0$ exist. Suppose that

$$F^{(m)}(0) = G^{(m)}(0) = 0,\ m = 0, 1, 2, \ldots, n-1 \ \text{ and } \ G^{(n)}(0) \neq 0.$$

Then we have

$$\lim_{x\to 0} \frac{F(x)}{G(x)} = \frac{F^{(n)}(0)}{G^{(n)}(0)}.$$

First of all we take $q = 0$. Then we have

$$\mathcal{D}_0 f(x) := \frac{f(x) - f(0)}{x},\ 0 < |x| < \rho$$

and $(\mathcal{D}_0 f)(0) := f'(0)$. Further we have

$$\begin{aligned}(\mathcal{D}_0^n f)(0) &:= \left(\mathcal{D}_0^{n-1} f\right)'(0) \\ &= \lim_{x\to 0} \frac{\mathcal{D}_0^{n-1} f(x) - \left(\mathcal{D}_0^{n-1} f\right)(0)}{x} = \lim_{x\to 0} \mathcal{D}_0^n f(x),\ n = 1, 2, 3, \ldots. \end{aligned} \tag{2}$$

Now we will show that

$$x^n \mathcal{D}_0^n f(x) = f(x) - \sum_{k=0}^{n-1} \frac{f^{(k)}(0)}{k!} x^k,\ 0 < |x| < \rho \tag{3}$$

and

$$(\mathcal{D}_0^n f)(0) = \frac{f^{(n)}(0)}{n!} \tag{4}$$

for each $n \in \{1, 2, 3, \ldots\}$ for which $f^{(n)}(0)$ exists.

We use induction on n. For $n = 1$ we have (3) and (4) by definition. So we assume that $f^{(n+1)}(0)$ exists and that (3) and (4) are valid for some $n \in \{1, 2, 3, \ldots\}$. Then we find

$$x^{n+1} \mathcal{D}_0^{n+1} f(x) := x^n \left[\mathcal{D}_0^n f(x) - (\mathcal{D}_0^n f)(0)\right]$$

$$
\begin{aligned}
&= f(x) - \sum_{k=0}^{n-1} \frac{f^{(k)}(0)}{k!} x^k - \frac{f^{(n)}(0)}{n!} x^n \\
&= f(x) - \sum_{k=0}^{n} \frac{f^{(k)}(0)}{k!} x^k,\ 0 < |x| < \rho.
\end{aligned}
$$

Further we have by using (2) and l'Hospital's rule above

$$
\begin{aligned}
\left(\mathcal{D}_0^{n+1} f\right)(0) &= \lim_{x \to 0} \mathcal{D}_0^{n+1} f(x) \\
&= \lim_{x \to 0} \frac{f(x) - \sum\limits_{k=0}^{n} \frac{f^{(k)}(0)}{k!} x^k}{x^{n+1}} = \frac{f^{(n+1)}(0)}{(n+1)!}.
\end{aligned}
$$

This proves (3) and (4) for each $n \in \{1, 2, 3, \ldots\}$ for which $f^{(n)}(0)$ exists.

Now a combination of (2) and (4) completes the proof of the theorem if $q = 0$.

For $q = -1$ the definition (1) leads to

$$
2x\mathcal{D}_{-1} f(x) = f(x) - f(-x),\ 0 < |x| < \rho
$$

and $\left(\mathcal{D}_{-1} f\right)(0) = f'(0)$. So we have by using l'Hospital's rule

$$
\lim_{x \to 0} \mathcal{D}_{-1} f(x) = \lim_{x \to 0} \frac{f(x) - f(-x)}{2x} = f'(0) = \left(\mathcal{D}_{-1} f\right)(0)
$$

if $f'(0)$ exists. This proves the theorem for $n = 1$.

Further we have

$$
4x^2 \mathcal{D}_{-1}^2 f(x) = f(x) - f(-x) + f(-x) - f(x) = 0,\ 0 < |x| < \rho.
$$

And by using l'Hospital's rule we obtain

$$
\begin{aligned}
\left(\mathcal{D}_{-1}^2 f\right)(0) &:= \left(\mathcal{D}_{-1} f\right)'(0) \\
&= \lim_{x \to 0} \frac{\mathcal{D}_{-1} f(x) - \left(\mathcal{D}_{-1} f\right)(0)}{x} = \lim_{x \to 0} \frac{f(x) - f(-x) - 2xf'(0)}{2x^2} = 0
\end{aligned}
$$

if $f''(0)$ exists. This implies that

$$
\mathcal{D}_{-1}^2 f(x) = 0,\ |x| < \rho.
$$

Hence

$$
\mathcal{D}_{-1}^n f(x) = 0,\ |x| < \rho,\ n = 2, 3, 4, \ldots.
$$

So we have for $n \geq 2$: if $f^{(n)}(0)$ exists, then $f''(0)$ exists and $\mathcal{D}_{-1}^n f(x) = 0$ for $|x| < \rho$. This proves the theorem for $q = -1$.

In the sequel we assume that $q \neq 0$ and $q \neq -1$.

We show first that for every positive integer n

$$\lim_{x\to 0} \mathcal{D}_q^n f(x) = \frac{f^{(n)}(0)}{n!} \frac{(q;q)_n}{(1-q)^n} \tag{5}$$

if $f^{(n)}(0)$ exists.

The definition (1) of the q-derivative operator $\mathcal{D}_q$ leads to

$$(1-q)x\mathcal{D}_q f(x) = f(x) - f(qx),\ 0 < |x| < \min(\rho, \frac{\rho}{|q|}).$$

If we multiply by $(1-q)x$ and replace f by $\mathcal{D}_q f$ we obtain

$$(1-q)^2 x^2 \mathcal{D}_q^2 f(x) = f(x) - (1+q^{-1})f(qx) + q^{-1}f(q^2x),\ 0 < |x| < \min(\rho, \frac{\rho}{|q|^2}).$$

Now we use induction on n to see that we have for $n \in \{1, 2, 3, \ldots\}$:

$$(1-q)^n x^n \mathcal{D}_q^n f(x) = \sum_{k=0}^{n} (-1)^k q^{-k(n-1)+\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q f(q^k x),\ 0 < |x| < \min(\rho, \frac{\rho}{|q|^n}) \tag{6}$$

where the q-binomial coefficient $\begin{bmatrix} n \\ k \end{bmatrix}_q$ is defined by

$$\begin{bmatrix} n \\ k \end{bmatrix}_q := \frac{(q;q)_n}{(q;q)_k (q;q)_{n-k}}.$$

Let n be a positive integer and assume that $f^{(n)}(0)$ exists.

Now we define

$$F(x) := \sum_{k=0}^{n} (-1)^k q^{-k(n-1)+\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q f(q^k x)$$

and

$$G(x) := (1-q)^n x^n.$$

Then we have

$$G^{(m)}(0) = 0,\ m = 0, 1, 2, \ldots, n-1 \ \text{ and } \ G^{(n)}(0) = (1-q)^n n! \neq 0$$

and

$$F^{(m)}(0) = f^{(m)}(0) \sum_{k=0}^{n} (-1)^k q^{k(m-n+1)+\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q,\ m = 0, 1, 2, \ldots, n.$$

Now we use (see for instance [2])

$$(a;q)_n = \sum_{k=0}^{n} q^{\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q (-a)^k$$

for q real with $|q| \neq 1$ and a an arbitrary complex number, to obtain

$$\sum_{k=0}^{n}(-1)^k q^{k(m-n+1)+\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q = (q^{m-n+1};q)_n,\ m = 0,1,2,\ldots. \tag{7}$$

Hence

$$\sum_{k=0}^{n}(-1)^k q^{k(m-n+1)+\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q = (q;q)_n \delta_{mn},\ m = 0,1,2,\ldots,n.$$

So we have

$$F^{(m)}(0) = 0,\ m = 0,1,2,\ldots,n-1 \ \text{ and } \ F^{(n)}(0) = (q;q)_n f^{(n)}(0).$$

Applying l'Hospital's rule above we obtain

$$\lim_{x\to 0} \mathcal{D}_q^n f(x) = \lim_{x\to 0} \frac{\sum\limits_{k=0}^{n}(-1)^k q^{-k(n-1)+\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q f(q^k x)}{(1-q)^n x^n} = \frac{f^{(n)}(0)}{n!}\frac{(q;q)_n}{(1-q)^n}.$$

This proves (5).

Now we prove that for every positive integer n

$$\left(\mathcal{D}_q^n f\right)(0) = \frac{f^{(n)}(0)}{n!}\frac{(q;q)_n}{(1-q)^n} \tag{8}$$

if $f^{(n)}(0)$ exists.

We use induction on n.

For $n = 1$ we have (8) by definition.

Assume that $f^{(n+1)}(0)$ exists and (8) holds for some $n \in \{1,2,3,\ldots\}$. Now we use the definition (1) and (6) to find

$$\begin{aligned} \left(\mathcal{D}_q^{n+1} f\right)(0) &:= \left(\mathcal{D}_q^n f\right)'(0) = \lim_{x\to 0} \frac{\mathcal{D}_q^n f(x) - \left(\mathcal{D}_q^n f\right)(0)}{x} \\ &= \lim_{x\to 0} \frac{\sum\limits_{k=0}^{n}(-1)^k q^{-k(n-1)+\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q f(q^k x) - \frac{f^{(n)}(0)}{n!}(q;q)_n x^n}{(1-q)^n x^{n+1}} =: L. \end{aligned}$$

Let

$$F(x) := \sum_{k=0}^{n}(-1)^k q^{-k(n-1)+\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix}_q f(q^k x) - \frac{f^{(n)}(0)}{n!}(q;q)_n x^n$$

and

$$G(x) := (1-q)^n x^{n+1}.$$

Then we have by using (7)

$$F^{(m)}(0) = G^{(m)}(0) = 0,\ m = 0,1,2,\ldots,n$$

and

$$F^{(n+1)}(0) = (q^2;q)_n f^{(n+1)}(0) \text{ and } G^{(n+1)}(0) = (1-q)^n (n+1)! \neq 0.$$

So we have by using l'Hospital's rule again

$$L = \frac{F^{(n+1)}(0)}{G^{(n+1)}(0)} = \frac{(q^2;q)_n f^{(n+1)}(0)}{(1-q)^n (n+1)!} = \frac{f^{(n+1)}(0)}{(n+1)!} \frac{(q;q)_{n+1}}{(1-q)^{n+1}}.$$

This proves (8) for every $n \in \{1,2,3,\ldots\}$ for which $f^{(n)}(0)$ exists.

The complex variable case.

Now let f be a function of a complex variable z.

If $f'(0)$ exists, then we have by definition $(\mathcal{D}_q f)(0) = f'(0)$. Further we have for $q \neq 0$

$$\begin{aligned} \lim_{z\to 0} \mathcal{D}_q f(z) &= \lim_{z\to 0} \frac{f(z) - f(qz)}{(1-q)z} \\ &= \lim_{z\to 0} \left(\frac{f(z) - f(0)}{(1-q)z} - q \frac{f(qz) - f(0)}{(1-q)qz} \right) = \frac{f'(0)}{1-q} - q \frac{f'(0)}{1-q} = f'(0). \end{aligned}$$

For $q = 0$ we simply obtain

$$\lim_{z\to 0} \mathcal{D}_0 f(z) = \lim_{z\to 0} \frac{f(z) - f(0)}{z} = f'(0).$$

If $f''(0)$ exists, then f' exists in a neighbourhood of $z = 0$ which means that f is analytic in a neighbourhood of $z = 0$. This implies that $f^{(m)}(0)$ exists for $m = 0, 1, 2, \ldots$ and we have

$$f(z) = \sum_{m=0}^{\infty} \frac{f^{(m)}(0)}{m!} z^m, \ |z| < \rho \text{ for some } \rho > 0.$$

Applying the definition (1) we find

$$\mathcal{D}_q f(z) = \sum_{m=1}^{\infty} \frac{f^{(m)}(0)}{m!} \frac{1-q^m}{1-q} z^{m-1}, \ |z| < \min(\rho, \frac{\rho}{|q|}),$$

where $\min(\rho, \frac{\rho}{|q|})$ should be replaced by ρ in the case that $q = 0$.

Hence $\mathcal{D}_q f(z)$ is analytic in a neighbourhood of $z = 0$. Now we use induction to find for $n \in \{1,2,3,\ldots\}$:

$$\mathcal{D}_q^n f(z) = \sum_{m=n}^{\infty} \frac{f^{(m)}(0)}{m!} \frac{(q^{m-n+1};q)_n}{(1-q)^n} z^{m-n}, \ |z| < \min(\rho, \frac{\rho}{|q|^n}),$$

where $\min(\rho, \frac{\rho}{|q|^n})$ should be replaced by ρ if $q = 0$.

Hence $\mathcal{D}_q^n f(z)$ is analytic in a neighbourhood of $z = 0$. So we easily obtain

$$\lim_{z\to 0} \mathcal{D}_q^n f(z) = \left(\mathcal{D}_q^n f\right)(0) = \frac{f^{(n)}(0)}{n!} \frac{(q;q)_n}{(1-q)^n}.$$

This completes the proof of the theorem in the complex variable case.

Remark 1. Since we defined the q-derivative operator for functions f which are only assumed to be differentiable at $x = 0$, the well-known limit

$$\lim_{q \to 1} \mathcal{D}_q f(x) = f'(x)$$

no longer holds in general, but only for those x where f is differentiable.

Remark 2. In the complex variable case the definition (1) can be extended to complex values of q and then the theorem still holds for complex values of $q \neq 1$.

References

[1] Ph. Franklin. *A treatise on advanced calculus.* John Wiley & Sons, New York, 1940.

[2] G. Gasper and M. Rahman. *Basic hypergeometric series.* Encyclopedia of Math. and its Appl. **35**, Cambridge University Press, 1990.

J. Koekoek
Menelaoslaan 4
5631 LN Eindhoven
The Netherlands

R. Koekoek
Delft University of Technology
Faculty of Technical Mathematics and Informatics
Mekelweg 4
2628 CD Delft
The Netherlands